INDEX

Page

Acetate, calcium, method used in determining soil acidity 322–329
Achorion Schönleinii, causal organism of favus 415–418
Acid—
amino, in normal and blighted spinach .. 385–404
hydrochloric, used in—
analyzing greensand 485
determining water-soluble nitrogen in spinach 397–404
hydrofluoric, used in analyzing greensand. 489
lactic, in alfalfa silag 581
nitric, used in experiments with plowsole .. 508
nitrohydrochloric, used in experiments with plowsole 507
phosphomolybdic, used in determining nitrogen in spinach 399–404
phosphotungstic, used in—
determining basic nitrogen in spinach. 399–404
titration of silage extract 130–131
sulphuric, used in—
damping-off experiment 536
determining basic nitrogen in spinach. 399–404
Acidity of—
alcoholic extract of silage 116–125
dioxogen 140–141
hydrogen peroxid 140–141
plant sap, changes during freezing 98–101
seeds, effect on catalase activity 141–143
silage 574, 581
soil, effect of moisture 321–329
water extract of silage 116–125
Acid-producers in alfalfa silage 574–592
Acids—
diamino, in healthy and blighted spinach. 399–404
tolerance of larvæ of *Trichinella spiralis* to. 477–479
Acromoniella sp., fungus isolated from *Ipomæa batatas* 364
Actinomyces sp., fungus isolated from *Ipomæa batatas* 364
Adams, J. M. R., et al. (paper): Sweet-Potato Storage-Rots 337–368
Adelphocoris rapidus, attacking clover 281
Adobe soil, inorganic colloids 516
Aegle marmelos, immunity to *Pseudomonas citri* 662–665
After-ripening of seeds, effect on—
catalase activity 138, 149–159
oxidase activity 163–166
Agriotes mancus, larvæ, effect of hydrocyanic-acid gas on 134–136
Agropyron—
coninum, source of *Puccinia graminis tritici* transferred to semicongenial hosts 242
cristatum—
source of *Puccinia graminis* in bridging host experiment 232
source of *Puccinia graminis tritici* transferred to semicongenial hosts 242

Agropyron—Continued Page
repens—
source of *Puccinia graminis* in bridging-host experiment 228
susceptibility to *Puccinia graminis tritici*. 224
smithii—
susceptibility to *Puccini graminis tritici* .. 224
used in bridging-host experiment 231–249
tenerum—
source of *Puccinia graminis tritici* transferred to semicongenial hosts 242
susceptibility to *Puccinia graminis tritici*. 224
Agrostis—
alba, source of *Puccinia graminis agrostis* transferred to uncongenial hosts 245
stolonifera—
source of *Puccinia graminis* in bridging-host experiment 228
source of *Puccinia graminis agrostis* transferred to uncongenial hosts 245
Albuminoid nitrogen in silage 129–132
Alcohol—
in silage—
caused by zymase 590
released by enzymic and yeast action .. 589–590
tolerance of larvæ of *Trichinella spiralis* to 477–479
Alcoholic extract of silage—
acidity 116–125
method of making 115
Alfalfa—
effect on plowsole 507
hay—
digestibility 271, 282
methane fermentation 274–275
net energy value 269–286
silage—
bacteria in 571–592
comparison with sweet-clover silage ... 113–132
Alkali—
effect of soil factors on toxity 287–319
salts, relative toxity 315–318
Alleys, effect on border rows in plot tests .. 251–262
Alisma spp., nature of dormancy of seeds 138
Alternaria sp., isolated from—
dead conifers 548
Ipomoea batatas 348, 355
Aluminum—
in mulch, plowsole, and subsoil 508–509
oxid in—
mulch, plowsole, and subsoil 508–516
normal and blighted spinach 371–375
soil, effect of drying 325–358
Amaranthus—
blitoides, catalase content of seeds 144, 153–154
graecizans, catalase content of seeds 153–154
retroflexus, catalase content of seeds 144, 149, 153–161
spp. nature of dormancy of seeds 138

Ambrosia— Page
artemisiaefolia, catalase and oxidase content of seeds ... 138
trifida, catalase and oxidase content of seeds 138
Amids, acid, in normal and blighted spinach ... 385–404
Amino—
acids in normal and blighted spinach ... 385–404
nitrogen in—
alfalfa silage ... 581–590
hardened and nonhardened cabbage ... 97
silage extract ... 125–132
Ammonia—
effect of carbon disulphid and toluol on accumulation in soil ... 603–614
in alfalfa silage ... 586–590
in hardened and nonhardened cabbage ... 97
Ammonia nitrogen in healthy and blighted spinach ... 398–404
Ammonium—
carbonate, used in experiments with plow-sole ... 508
hydrate, used in experiments with plow-sole ... 508
Amygdalus persica, nature of dormancy of seeds ... 138
Andropogon halepensis sudanensis, catalase and oxidase content of seeds ... 137–174
Angular-Leafspot of Cucumber: Dissemination, Overwintering, and Control (paper). 201–220
Angular-leafspot of cucumber—
dissemination by—
drainage water ... 211
insects ... 212–213
pickers ... 211–212
rain and wind ... 208–210
effect of—
desiccation ... 203–204
germicides ... 205
fruit infection ... 208
host plants ... 205–206
stomatal infection ... 206–208
thermal death point ... 204–205
overwintering in—
insects ... 214
seeds ... 214
soil ... 213
remedial measures—
resistant varieties ... 217
sanitation ... 217
seed treatment ... 218
spraying ... 217–218
Ant. See *Pheidole megacephala*.
Armsby, Henry Prentiss, and Fries, J. August (paper): Net Energy Values of Alfalfa Hay and of Starch ... 269–286
Arny, A. C., and Hayes, H. K. (paper): Experiments in Field Technic in Plot Tests. 251–262
Artschwager, Ernst F. (paper): Histological Studies on Potato Leafroll ... 559–570
Ash—
digestibility in alfalfa hay and starch ... 271, 282
in normal and blighted spinach ... 371–375
in dry matter of feeding stuffs ... 270
Asparagin, used in determining water-soluble nitrogen in spinach ... 397–404
Aspergillus— Page
niger, fungus isolated from *Ipomoea batatas*. 364
sp., not cause of damping-off in conifers ... 549
Allantia—
cirtrioides, susceptibility to *Pseudomonas citri* ... 662–665
disticha, susceptibility to *Pseudomonas citri* ... 662–665
Aucuba japonica, production of injected areas due to freezing ... 86
Austrian pine. See *Pinus austriaca*.
Avena sativa—
reported bridging host for transfer of timothy rust ... 222
used in bridging-host experiments ... 226–249
Azotobacter, effect of carbon disulphid and toluol on ... 601–614
Bacteria, nitrogen-fixing, effect of carbon disulphid and toluol ... 601–614
Bacteriological Studies on Alfalfa Silage (paper) ... 571–592
Bacterium lachrymans, causal organism of angular-leafspot of cucumber ... 201–220
Bactrocera cucurbitae—
host of *Opius fletcheri* ... 456–457
host of *Tetrastichus giffardianus* ... 454–458
Balsam-apple. See *Momordica balsamina*.
Balsamocitrus gabonensis, immunity to *Pseudomonas citri* ... 662–665
Balsam-pear. See *Momordica charantia*.
Barberry. See *Berberis* spp.
Barium—
chlorid, used in determining nitrogen in spinach ... 400–404
hydroxid, used in determining nitrogen in spinach ... 399–404
phosphotungstate, used in determining nitrogen in spinach ... 399–404
Barley. See *Hordeum* spp.
Basic nitrogen in healthy and blighted spinach ... 399–404
Basswood. See *Tilia americana*.
Beach, B. A, and Halpin, J. G. (paper): Observations on an Outbreak of Favus ... 415–418
Beef cattle, crosses with dairy breeds ... 1–57
Beetle, June green. See *Colinus nitida*.
Begonia, production of injected areas due to freezing ... 86
Berberis spp., reported bridging host for *Puccina poculiformis* ... 222
Bermuda grass. See *Capriola dactylon*.
Bestill. See *Thevetia neriifolia*.
Beta vulgaris, effect of farm manure on ... 494–503
Biologic forms of *Puccinia graminis*, plasticity of ... 221–249
Biology of fruit-fly parasites in Hawaii ... 419–466
Black, Otis F., et al. (paper): Physiological Studies of Normal and Blighted Spinach. 369–408
Blackrot—
Java, of *Ipomoea batatas* ... 347–349
of *Ipomoea batatas*—
description ... 344–346
susceptible varieties ... 346
Blacktop, type of damping-off ... 525–558
Black-walnut worm. See *Datana integerrima*.
Bluegrass. See *Poa pratensis*.

Page

Bordeaux mixture, used against *Bacterium lachrymans*... 217-218
Botryorhiza hippocrateae, difference from *Cronartium ribicola*... 627-628
Botrytis—
  cinerea—
    cause of damping-off in conifers... 547-558
    fungus isolated from *Ipomoea batatas*... 356-357
  douglasii. Syn. *Botrytis cinerea*.
  vulgaris. Syn. *Botrytis cinerea*.
Bouyoycos, George J., and McCool, M. M. (paper): Determining the Absolute Salt Content of Soils by Means of the Freezing-Point Method... 331-336
Brassica—
  napus, injected areas due to freezing... 87
  oleracea capitata, frost injury in... 85
Brier roses, resistant to brown canker... 593-600
Brisket disease—
  condition of heart... 409-411
  effect of—
    altitude... 411-414
    feed... 411-413
Bromelin, effect on oxidase activity of seeds... 169
Bromus tectorum—
  susceptibility to *Puccinia graminis*... 222, 228-229, 240-241
  used in bridging-host experiment... 228-249
Brown Canker of Roses, Caused by Diaporthe umbrina (paper)... 593-600
Brown rust of bromes. See *Puccinia dispersa*.
Brown-tail moth. See *Euproctis chrysorrhoea*.
Bryanopsis laciniosa, host plant of *Bacterium lachrymans*... 206
Bug, gray-damsel. See *Reduviolus ferus*.
Bunzell, H. H., et al. (paper): Physiological Studies of Normal and Blighted Spinach. 369-408
Bulgarian bacteria in alfalfa silage... 581-590
Byrophyllum, production of injected areas due to freezing... 86-88
Cabbage. See *Brassica oleracea capitata*.
Calabash gourd. See *Lagenaria vulgaris*.
Calcium—
  carbonate—
    effect on plowsole... 507
    in olive earth... 484
    used to neutralize hydrogen peroxid... 142-143
  in mulch, plowsole, and subsoil... 508-516
  malophosphate crystals, formation in frozen spots of cabbage leaf... 86-87
  nitrate in solution used in experiments with greensand... 486-492
  oxid in—
    greensand... 485
    mulch, plowsole, and subsoil... 515-516
    normal and blighted spinach... 371-375
    soil, effect of drying... 325-328
    sulphate, effect on plowsole... 507
Calcium-acetate method, used in determining soil acidity... 322-329
Cane sugar, effect on alfalfa silage... 584-590
Canker, brown, of roses... 593-600
Capnodium meridionale, similarity to *Coleosporium senecionis*... 648
Capriola dactylon, catalase and oxidase content of seeds... 137-174

Page

Carbohydrate changes during hardening against frost injury... 95-97
Carbohydrate production in healthy and blighted spinach... 381-384
Carbohydrates, effect on alfalfa silage... 584-590
Carbon—
  digestibility in alfalfa hay and starch... 271, 282
  dioxid in—
    gaseous excretion... 273, 284
    silage, released by enzymic and yeast action... 589-590
  disulphid, effect on nitrogen-fixing organisms in soil... 601-614
  in dry matter of feeding stuffs... 270
  in gaseous excretion... 273, 284
Carbonate—
  ammonium, used in experiments with plowsole... 508
  calcium—
    effect on plowsole... 507
    in olive earth... 484
  sodium, effect on plowsole... 507
Carsner, Eubanks (paper): Angular-Leafspot of Cucumber: Dissemination, Overwintering, and Control... 201-220
Caryopsis of seeds, catalase activity of... 146-147
Casein digesters in alfalfa silage... 586-590
Catalase and Oxidase Content of Seeds in Relation to Their Dormacy, Age, Vitality, and Respiration (paper)... 137-174
Catalase of seeds—
  method of determining... 139-141
  solubility... 160-161
Cattle, crosses between dairy and beef breeds... 1-37
Cattleya orchids, fumigation with hydrocyanic-acid gas... 263-268
Cells, diameter in various parts of seeds of Johnson grass and crimson clover... 145
Cephalothecium sp., fungus isolated from *Ipomoea batatas*... 364
Ceratitis capitata in Hawaii, parasitism... 419-466
Ceratocystis fimbriata. See *Sphaeronema fimbriatum*.
Ceratostoma sp., fungus isolated from *Ipomoea batatas*... 364
Chaetomium sp., not cause of damping-off in conifers... 540-558
Chaetospermum glutinosa, susceptibility to *Pseudomonas citri*... 662-665
Chalcas exotica, susceptibility to *Pseudomonas citri*... 661-666
Charcoal-rot of *Ipomoea batatas*... 351-352
Chemical changes in plants during hardening against frost injury... 95-97
Chemistry of Sweet-Clover Silage in Comparison with Alfalfa Silage (paper)... 113-132
Cherry, French. See *Eugenia uniflora*.
Chlorid—
  barium, used in determining nitrogen in spinach... 400-404
  mercuric—
    effect on *Bacterium lachrymans*... 205
    effect on oxidase activity of seeds... 168-169
    used in determining nitrogen in spinach... 399-404

Chlorid—Continued. Page
potassium, used in experiments with greensand ... 488
silver, used in determining nitrogen in spinach ... 400-404
sodium—
effect on plowsole ... 507
used in alkali experiment ... 289-319
used in experiments with intestinal trichinæ ... 468-482
Chloroform, effect on alfalfa silage ... 590
Chrysophyllum monopyrenum, host of puparia parasitized by *Diachasma fullawayi* ... 448
Cimex—
antenni-rectus. See *Miris dolabratus*.
deses. See *Miris dolabratus*.
dolabratus. See *Miris dolabratus*.
frumentaruis. See *Miris dolabratus*.
laevigatus. See *Miris dolabratus*.
lateralis. See *Miris dolabratus*.
porrectus. See *Miris dolabratus*.
recticornis. See *Miris dolabratus*.
riparius. See *Miris dolabratus*.
V-flavum. See *Miris dolabratus*.
Citratae, susceptibility to *Pseudomonas citri* 662-665
Citreae, susceptibility to *Pseudomonas citri*. 662-665
Citrinae, susceptibility to *Pseudomonas citri* 662-665
Citron. See *Citrullus vulgaris*.
Citrullus vulgaris—
free from attacks of *Bacterium lachrymans* ... 206
host of organism virulent to *Ipomoea batatas* 347
Citrus canker. See *Pseudomonas citri*.
Citrus—
grandis, susceptibility to *Pseudomonas citri* ... 664-665
sinensis, susceptibility to *Pseudomonas citri* ... 665
Citropsis schweinfurthii, susceptibility to *Pseudomonas citri* ... 662-665
Clays, potash in ... 70-72
Claucena lansium, susceptibility to *Pseudomonas citri* ... 662-665
Clover—
crimson. See *Trifolium incarnatum*.
red. See *Trifolium pratense*.
Coffea arabica, host of puparia parasitized by *Diachasma fullawayi* ... 448
Coffee. See *Coffea arabica*.
Coleoptera, found on imported orchids ... 263
Coleosporium—
senecionis, similarity to *Caeoma meridionale* ... 648
solidaginis. Syn. *Coleosporium sonchi-arvensis*.
sonchi-arvensis, fusion nucleus in teliospore. 646
Coleus, production of injected areas due to freezing ... 86
Collembola, found on imported orchids ... 263
Colley, Reginald H. (paper): Parasitism, Morphology, and Cytology of Cronartium ribicola ... 619-660
Colloids, soil, relation to plowsoles ... 505-519
Colocasia esculenta, host of rot-producing organism virulent to *Ipomoea batatas* ... 347
Colon organisms in alfalfa silage ... 574-592
Colorimetric titration of silage extract ... 127-129
Complement-fixation test, serum distributed by multiple-pipette holder ... 615-618
Condition of Fertilizer Potash Residues in Hagerstown Silty Loam Soil (paper) ... 59-81
Conifers, diseases of seedlings ... 521-558
Conifer seedlings, injury from—
damping-off ... 521-558
drouth ... 554-555
mechanical means ... 551-554
whitespot ... 551
Conner, S. D. (paper): Soil Acidity as Affected by Moisture Conditions of the Soil ... 321-329
Contribution to the Biology of Fruit-Fly Parasites in Hawaii (paper) ... 419-466
Copper—
sulphate, effect on *Bacterium lachrymans* ... 205
sulphid, used in determining nitrogen in spinach ... 400-404
Coriseus ferus, natural enemy of *Leptoterma dolobrata* ... 194
Corn chop, effect on alfalfa silage ... 574-590
Corrodentia, found on imported orchids ... 263
Corsican pine. See *Pinus nigra poiretiana*.
Corticium vagum, cause of damping-off in conifers ... 522-558
Cotinus nitida, larvæ, effect of hydrocyanic-acid gas on ... 134-136
Crataegus spp., nature of dormancy of seeds ... 138
Crimson clover. See *Trifolium incarnatum*.
Crocker, William, and Harrington, George T. (paper): Catalase and Oxidase Content of Seeds in Relation to Their Dormancy, Age, Vitality, and Respiration ... 137-174
Cronartium—
ascelpiadeum, resemblance to *Cronartium ribicola* ... 653
comptoniae, nature of injury on *Pinus sylvestris* ... 649
flaccidum, parasite on *Peonia officinalis* ... 638-639
occidentale, æcia ... 652
ribicola—
cytology ... 639-653
infection of pine ... 624-625
life history ... 620-624
morphology—
æcia ... 630-634
mycelium ... 626-629
pycnia ... 629-630
telia ... 637-639
uredinia ... 635-637
Crookneck squash, Japanese. See *Cucurbita moschata*.
Crosses between dairy and beef cattle—
breeds in parent herd ... 2
inheritance of—
beef qualities ... 49-51
color ... 22-24
milking qualities ... 51-52
muzzle color ... 42-43
polled character ... 45-48
switch color ... 44
tongue pigment ... 43-44
white markings ... 24-42
Crude fiber—
digestibility in alfalfa hay and starch ... 271, 282
in dry matter of feeding stuffs ... 270

Cucumber— Page
squirting. See *Momordica elaterium*.
wild. See *Echinocystis lobata*.
See *Cucumis sativus*.
Cucumis—
aculangulis, host plant of *Bacterium lachrymans* 206
anguria, host plant of *Bacterium lachrymans* 206
dipsaceus, host plant of *Bacterium lachrymans* 206
grossulariaeformis, free from attacks of *Bacterium lachrymans* 206
melo—
free from attacks of *Bacterium lachrymans* 206
var. *dudaim*, free from attacks of *Bacterium lachrymans* 206
var. *flexuosus*, free from attacks of *Bacterium lachrymans* 206
sativus, host plant of *Bacterium lachrymans*. 206
Cucurbita—
maxima, free from attacks of *Bacterium lachrymans* 206
moschata, free from attacks of *Bacterium lachrymans* 206
pepo—
free from attacks of *Bacterium lachrymans* 206
var. *condensa*, free from attacks of *Bacterium lachrymans* 206
var. *ovifera*, free from attacks of *Bacterium lachrymans* 206
Cuprous oxid in intumescences on frozen leaves 88
Cyanid—
potassium, tolerance of larvæ of *Trichinella spiralis* to 477–479
sodium—
effect on cattleya orchids 267–268
used to generate hydrocyanic-acid gas 134
Dactylis glomerata—
food plant of *Miris dolabratus* 182
reported bridging host for transfer of timothy rust 222
source of *Puccinia graminis phleipratensis* transferred to semicongenial hosts 244
Dairy cattle, crosses with beef breeds 1–57
Damping-off diseases of conifers 521–558
Dasheen. See *Colocasia esculenta*.
Datana integerrima, effect of hydrocyanic-acid gas on 264–268
Denitrification in diseased spinach 401–404
Desiccation, effect on catalase activity of seeds 149–150
Determining the Absolute Salt Content of Soils by Means of the Freezing-Point Method (paper) 331–336
Dextrose in intumescences on frozen leaves 88
Diabrotica—
duodecimpunctata, possible carrier of *Bacterium lachrymans* 212
vittata, possible carrier of *Bacterium lachrymans* 212
Diachasma—
fullawayi—
life history 445–448
natural enemy of Mediterranean fruit fly 419–466

Diachasma—Continued. Page
tryoni—
life history 420–439
natural enemy of Mediterranean fruit fly 419–466
Diamino acids in healthy and blighted spinach 399–404
Diaporthe—
batatatis, causal organism of dryrot of *Ipomoea batatas* 349
umbrina, n. sp., cause of brown canker of roses 593–600
control 598–599
cultural characters 598
description 596–597
life history 598
Dicotyledons, parasitized by same damping-off parasite as conifers 535–558
Dietz, H. F., and Sasscer, E. R. (paper): Fumigation of Cattleya Orchids with Hydrocyanic-Acid Gas 263–268
Dimorphism of *Miris dolabratus* 183–184
Dioxid—
carbon, in silage, released by enzymic and yeast action 589–590
silicon in—
mulch, plowsole, and subsoil 508–516
normal and blighted spinach 371–375
Dioxygen, used in determining catalase content of seeds 140–144
Diplodia—
gossypina, causal organism of rot in *Ipomoea batatas* 347
macrura, causal organism of rot in *Ipomoea batatas* 347
natalensis, causal organism of rot in *Ipomoea batatas* 347
tubericola, causal organism of Java blackrot of *Ipomoea batatas* 347
zeae, nonvirulent to *Ipomoea batatas* 348
Disaccharids in hardened and nonhardened cabbage 95–96
Discosia pini, not cause of damping-off in conifers 550
Diseases of conifer seedlings 521–558
Disulphid, carbon, effect on nitrogen-fixing organisms in soil 601–614
Dormancy of seeds, relation of catalase and oxidase content to 137–174
Douglas fir. See *Pseudotsuga taxifolia*.
Drouth injury, compared with damping-off in conifer seedlings 554–555
Drying, effect on catalase activity of seeds 149–150
Dryrot of *Ipomoea batatas* 349–350
Echinocystis lobata, free from attacks of *Bacterium lachrymans* 206
Effect of Carbon Disulphid and Toluol upon Nitrogen-Fixing and Nitrifying Organisms (paper) 601–614
Effect of Farm Manure in Stimulating the Yields of Irrigated Field Crops (paper) 493–503
Effect of Hydrocyanic-Acid Gas under Vacuum Conditions on Subterranean Larvæ (paper) 133–136
Elaeagnus sp., host of damping-off parasites, 535–558
Electrometric titration of silage extract 125–127

Elymus— Page
canadensis, used in bridging-host experiments ... 226–249
macounii, source of *Puccinia graminis tritici* transferred to semicongenial hosts ... 242
robustus used in bridging-host experiment, ... 234–249
virginicus, used in bridging-host experiment ... 233–249
Embryo of seeds, catalase activity of ... 146–147
Endophyllum sempervivi, æciospore chains ... 647
Endosperm of seeds, catalase activity of ... 146–147
Energy expenditure consequent upon feed consumption ... 278–280
Engelmann spruce. See *Picea engelmanni*.
Entomophthora sp., fungus attacking *Miris dolabratus* ... 197
Enzyms in alfalfa silage ... 589–590
Epicoccum sp., fungus isolated from *Ipomoea batatas* ... 357
Erb, E. S., and Frear, William (paper): Condition of Fertilizer Potash Residues in Hagerstown Silty Loam Soil ... 59–81
Eremocitrus glauca, susceptibility to *Pseudomonas citri* ... 662–665
Eriobotrya japonica, host fruit of puparia parasitized by *Diachasma fullawayi* ... 448
Errata and authors' emendations ... IV
Erysiphe graminis, tested on *Bromus* spp. ... 222
Ether extract in dry matter of feeding stuffs ... 270
Ethyl-acetate method, used in determining soil acidity ... 322–329
Eugenia uniflora—
host fruit of puparia parasitized by *Diachasma fullawayi* ... 448
host plant of puparia parasitized by *Diachasma tryoni* ... 429
Eugenol, used to determine oxidase activity in normal and blighted spinach ... 377–380
Eulophinae, one species natural enemy of Mediterranean fruit fly ... 419–466
Euproctis chrysorrhoea, used in experiments with *Limnerium validum* ... 458
Evodia—
latifolia, susceptibility to *Pseudomonas citri* ... 663–665
ridleyei, susceptibility to *Pseudomonas citri* ... 663–665
Excretion—
gaseous, in alfalfa hay and starch-feeding experiment ... 273, 284
urinary, in alfalfa hay and starch-feeding experiment ... 273, 283
Experiments in Field Technic in Plot Tests (paper) ... 251–262
Experiments on the Value of Greensand as a Source of Potassium for Plant Culture (paper) ... 483–492
Fall webworm. See *Hyphantria cunea*.
Favus, skin disease caused by *Achorion Schönleinii*—
anatomical changes ... 416
enzootiology ... 416–417
treatment ... 417–418
Feed—
effect on brisket disease ... 411–414
influence of quantity on digestibility ... 272
relation to energy expenditure ... 278–280
Feeding stuffs, composition and energy content of ... 270
Fermentation, methane, in alfalfa hay and starch ... 274–275
Feroniella lucida, susceptibility to *Pseudomonas citri* ... 662–665
Feronia limonia, susceptibility to *Pseudomonas citri* ... 662–665
Feroninae, susceptibility to *Pseudomonas citri* ... 662–665
Ferric oxid in—
mulch, plowsole, and subsoil ... 508–516
soil, effect of drying ... 325–328
Fertilizer potash residues in Hagerstown silty loam soil ... 59–81
Fescue, meadow. See *Festuca elatior*.
Festuca—
elatior—
food plant of *Miris dolabratus* ... 182
reported bridging host for transfer of timothy rust ... 222
source of *Puccinia graminis phleipratensis* transferred to semicongenial hosts ... 244
pratensis, source of *Puccinia graminis phleipratensis* transferred to semicongenial hosts ... 244
Fir, Douglas. See *Pseudotsuga taxifolia*.
Fly, fruit. See *Ceratitis capitata*.
Footroot of *Ipomoea batatas* ... 350
Forests, damping-off in ... 525–558
Formaldehyde—
effect on *Bacterium lachrymans* ... 205
remedy for favus ... 417–418
Formol, used in determining nitrogen in spinach ... 400–404
Fortunella—
hindsii, susceptibility to *Pseudomonas citri* ... 662–665
japonica, susceptibility to *Pseudomonas citri* ... 662–665
Four-lined plant bug. See *Poecilocapsus lineatus*.
Frost injury to plants, growth in injected areas ... 87–88
Frear, William, and Erb, E. S. (paper): Condition of Fertilizer Potash Residues in Hagerstown Silty Loam Soil ... 59–81
Freezing in plants ... 83–111
Freezing-point method, determination of salt content of soil by ... 331–336
Freezing points of juice from hardened and nonhardened plants ... 94–95
French cherry. See *Eugenia uniflora*.
Fries, J. August, and Armsby, Henry Prentiss (paper): Net Energy Values of Alfalfa Hay and of Starch ... 269–286
Frost injury—
first indications in cabbage and tomato ... 85
hardening cabbage against—
chemical changes ... 95–97
physiological changes ... 92–93
peroxidase changes in injured areas ... 88–90
"Frosted grains," relation to after-ripening ... 153
Fruit fly. See *Ceratitis capitata*.
Fumigation of Cattleya Orchids with Hydrocyanic-Acid Gas (paper) ... 263–268

Page

Fumigation with hydrocyanic-acid gas, effect on subterranean larvæ..... 133-136
Fungous parasites, cause of damping-off in conifers..... 521-558
Further Data on the Susceptibility of Rutaceous Plants to Citrus-Canker (paper).... 661-666
Further Studies on Brisket Disease (paper). 409-414
Fusarium—
acuminatum—
fungus used in inoculation experiments with *Ipomoea batatas*..... 360
not cause of damping-off in conifers.... 540-558
blasticola, cause of damping-off in conifers. 537-546
caudatum, fungus isolated from *Ipomoea batatas*..... 363-364
culmorum, fungus used in inoculation experiments with *Ipomoea batatas*..... 359-360
hyperoxysporum, fungus isolated from *Ipomoea batatas*..... 363-364
moniliforme, cause of damping-off in conifers..... 522-558
orthoceras, fungus isolated from *Ipomoea batatas*..... 363-364
oxysporum—
fungus isolated from *Ipomoea batatas*..... 360
isolated in sweet-potato storage-rots..... 338, 360, 363-364
parasiticum, cause of damping-off in conifers..... 537-546
pini, cause of damping-off in conifers..... 537-546
solani, cause of damping-off in conifers... 540-558
vasinfectum—
cause of damping-off in conifers..... 537-546
fungus isolated from *Ipomoea batatas*..... 364
ventricosum, cause of damping-off in conifers..... 522-558
Fusoma spp., reported cause of damping-off in conifers..... 537-546
Gainey, P. L. (paper): Effect of Carbon Disulphid and Toluol upon Nitrogen-Fixing and Nitrifying Organisms..... 601-614
Galesus silvestrii, parasite of Mediterranean fruit fly..... 433-466
Gas, hydrocyanic-acid—
effect on subterranean larvæ..... 133-136
fumigation of cattleya orchids..... 263-268
Gaseous excretion in alfalfa hay and starch feeding experiment..... 273, 284
Gastric digestion, effect on larvæ of *Trichinella spiralis*..... 470-471
Geise, Fred W., and True, Rodney H. (paper): Experiments on the value of Greensand as a Source of Potassium for Plant Culture... 483-492
Gelatin liquefiers in alfalfa silage..... 581-590
Geranium, production of injected areas due to freezing..... 86
Germinator, effect on catalase activity of seeds 156-159
Germination, effect on oxidase activity of seeds..... 166-167
Gherkin, West Indian. *Cucumis anguria*.
Gibberella saubinetii, fungus used in inoculation experiments with *Ipomoea batatas*... 357-358
Glauconite, applied to farm crops..... 483-492
Glover, George H., and Newsom, L. E. (paper): Further Studies on Brisket Disease. 409-414

Glucose in— Page
hardened and nonhardened cabbage..... 95
normal and blighted spinach..... 383
Glycerin, used in determining carbohydrates in normal and blighted spinach..... 383
Glycosmis pentaphylla, immunity to *Pseudomonas citri*..... 664-665
Gourd—
calabash. See *Lagenaria vulgaris*.
hedgehog. See *Cucumis dipsaceus*.
mandera. See *Cucumis acutangulis*.
See *Cucurbita pepo* var. *ovifera*.
Gowen, John W. (paper): Studies in Inheritance of Certain Characters of Crosses between Dairy and Beef Breeds of Cattle.... 1-38
Grass—
Bermuda. See *Capriola dactylon*.
Johnson. See *Sorghum halepensis*.
orchard. See *Dactylis glomerata*.
Sudan. See *Andropogon halepensis sudanensis*.
Tunis-sorghum hybrid, oxidase activity. 161-169
witch. See *Panicum capillare*.
Gray-damsel bug. See *Reduviolus ferus*.
Green June beetle. See *Cotinus nitida*.
Greensand, source of potassium..... 483-492
Grub, white. See *Lachnosterna* sp.
Guaiacol, used to determine oxidase activity in normal and blighted spinach..... 377-380
Guinea pig, weight of heart..... 411
Gypsum, used in experiments with plow-sole..... 514-515
Gypsy moth. See *Porthetria dispar*.
Halpin, J. G., and Beach, B. A. (paper): Observations on an Outbreak of Favus..... 415-418
Halticus citri, pest of beans..... 181
Hardening Process in Plants and Developments from Frost Injury (paper)..... 83-111
Hardiness to frost, method of testing..... 93
Hardiness to frost, time and temperature factors in acquiring or losing..... 93-94
Harrington, George T., and Crocker, William (paper): Catalase and Oxidase Content of Seeds in Relation to Their Dormancy, Age, Vitality, and Respiration..... 137-174
Harris, F. S., and Pittman, D. W. (paper): Soil Factors Affecting the Toxicity of Alkali 287-319
Harter, L. L., Weimer, J. L., and Adams, J. M. R. (paper): Sweet-Potato Storage-Rots 337-368
Hartley, Carl, and Merrill, T. C. (paper): Seedling Diseases of Conifers..... 521-558
Harvey, R. B. (paper): Hardening Process in Plants and Developments from Frost Injury..... 83-111
Hawaii, fruit-fly parasites in..... 419-466
Hawkins, Lon A., et al. (paper): Physiological Studies of Normal and Blighted Spinach..... 369-408
Hay, alfalfa—
digestibility..... 271, 282
net energy value..... 269-286
Hayes, H. K., and Army, A. C. (paper): Experiments in Field Technic in Plot Tests..... 251-262

Heart— Page
condition in brisket disease ... 409-411
weight in—
guinea pig ... 411
rabbit ... 411
Heat production and emission in feeding experiment ... 277-278, 284-286
Hedgehog gourd. See *Cucumis dipsaceus.*
Helianthus annuus, production of injected areas due to freezing ... 86
Hemiptera, found on imported orchids ... 263
Hesperethusa crenulata, susceptibility to *Pseudomonas citri* ... 662-665
Histological Studies on Potato Leafroll (paper) ... 559-570
Holcus halepensis sudanensis. Syn. *Andropogon halepensis sudanensis* and *Sorghum halepensis.*
Hordeum—
jubatum—
source of *Puccinia graminis tritici* transferred to semicongenial hosts ... 242
source of *Puccinia graminis* in bridging-host experiment ... 230
pusillum, used in bridging-host experiment ... 230-249
spp., used in bridging-host experiments ... 223-249
vulgare—
reported infected with timothy rust through bridging host ... 222
source of *Puccinia graminis tritici* transferred to semicongenial hosts ... 242
Hubbard squash. See *Cucurbita maxima.*
Humin nitrogen in healthy and blighted spinach ... 399-404
Humus extracts of mulch, plowsole, and subsoil, analysis ... 508
Hunter, O. W. (paper): Bacteriological Studies on Alfalfa Silage ... 571-592
Hydrangea, production of injected areas due to freezing ... 85
Hydrate, ammonium, used in experiments with plowsole ... 508
Hydrochloric acid. See Acid, hydrochloric.
Hydrocyanic-acid gas—
effect on subterranean larvæ ... 133-136
fumigation of cattleya orchids ... 263-268
generation of ... 134
Hydrofluoric acid. See Acid, hydrofluoric.
Hydrogen—
electrode, used in titration of silage extract 118-127
in gaseous excretion ... 273, 284
peroxid, used in determining catalase content of seeds ... 140-144
Hydrogen-ion concentration of—
plant sap, changes in during freezing ... 98-101
silage extract ... 118-127
Hydroquinone—
oxidase of Johnson grass not active toward ... 162
test for oxidase in—
frozen cabbage leaves ... 88
normal and blighted spinach ... 377-380
Hydroxid—
barium, used in determining nitrogen in spinach ... 399-404
sodium—
in alfalfa silage ... 586-590
Hydroxid—Continued. Page
sodium—Continued.
used to—
determine catalase activity of seeds ... 139
determine water-soluble nitrogen in spinach ... 398-404
neutralize hydrogen peroxid ... 142-144
Hymenoptera, found on imported orchids ... 263
Hyphantria cunea, host of *Limnerium validum.* 458
Hypochnus solani. Syn. *Corticium vagum.*
Hystrix patula, source of *Puccinia graminis* in bridging-host experiment ... 233
Inheritance of characters of crosses between dairy and beef cattle ... 1-57
Intestinal trichinæ. See *Trichinella spiralis.*
Ipomoea batatas—
storage-rots ... 337-368
varieties susceptible to blackrot ... 346
Irish potatoes. See *Solanum tuberosum.*
Iron in—
greensand ... 483-492
mulch, plowsole, and subsoil ... 508-516
normal and blighted spinach ... 371-375
Isoeugenol, used to determine oxidase activity in normal and blighted spinach ... 377-380
Jack pine. See *Pinus banksiana.*
Japanese crookneck squash. See *Cucurbita moschata.*
Java blackrot of *Ipomoea batatas—*
description ... 347-348
inoculation experiments ... 348-349
Jenkins, Anna E. (paper): Brown Canker of Roses, Caused by Diaporthe umbrina ... 593-600
Jodidi, Samuel L., et. al. (paper): Physiological Studies of Normal and Blighted Spinach ... 369-408
Johnson grass. See *Sorghum halepensis.*
June beetle, green. See *Cotinus nitida.*
Kamani nuts. See *Terminalia catappa.*
Kellogg, Edward H., et al. (paper): Physiological Studies of Normal and Blighted Spinach ... 369-408
Kelly, James W., et. al. (paper): Physiological Studies of Normal and Blighted Spinach ... 369-408
Kjeldahl method used in determination of silage extract ... 129-131
Lachnosterna sp., larvæ, effect of hydrocyanic-acid gas on ... 134-136
Lactic acid. See Acid, lactic.
Lagenaria vulgaris, host plant of *Bacterium lachrymans* ... 206
Larvæ, effect of hydrocyanic-acid gas on ... 133-136
Lasiodiplodia nigra, causal organism of rot in *Ipomoea batatas* ... 347
Latuca sativa, production of injected areas due to freezing ... 86
Lavanginae, susceptibility to *Pseudomonas citri* ... 662-665
Leafroll of *Solanum tuberosum* ... 559-570
Leafspot, angular, of cucumber ... 201-220
Lee, H. Atherton (paper): Further data on the Susceptibility of Rutaceous Plants to Citrus-Canker ... 661-666
Lepidoptera, found on imported orchids ... 263
Leptoterna—
dolabrata. See *Miris dolabratus.*
id. See *Miris dolabratus.*

Lettuce. See *Latuca sativa*. Page
Levine, M. N., et al. (paper): Plasticity of Biologic Forms of Puccinia graminis..... 221–249
Levulose in intumescences on frozen leaves .. 88
Lime in greensand 483–492
Limnerium validum, natural parasite of *Hyphantria cunea*..... 458
Loblolly pine. See *Pinus taeda*.
Lopomorphus id. See *Miris dolabratus*.
Lopus id. See *Miris dolabratus*.
Loquat. See *Eriobotrya japonica*.
Lycopersicon esculentum, first indications of frost injury in..... 85
Lygus pratensis, general plant pest..... 181
Macrosporium sp., fungus isolated from *Ipomoea batatas*..... 364
Magnesia, used in determining water-soluble nitrogen in spinach..... 398–404
Magnesium—
in mulch, plowsole, and subsoil 508–516
oxid—
in soil, effect of drying..... 325–328
in normal and blighted spinach..... 371–375
used in determining water-soluble nitrogen in spinach..... 398–404
sulphate—
effect on plowsole..... 507
in solution used in experiments with greensand..... 486–492
Mandera gourd. See *Cucumis dipsaceus*.
Manganese—
in plant tissues, relation to oxidase activity 373
oxid—
in soil, effect of drying..... 325–328
red, in normal and blighted spinach... 371–375
Manganous oxid in mulch, plowsole, and subsoil..... 515–516
Mangifera indica, host of rot-producing organism virulent to *Ipomoea batatas*..... 347
Mango. See *Mangifera indica*.
Manure, effect on irrigated crops..... 493–503
Marl, applied to farm crops..... 483–492
Maturity of seeds, effect on—
catalase activity..... 148–149
oxidase activity..... 163–164
Meadow fescue. See *Festuca elatior*.
Meadow Plant Bug, Miris dolabratus, The (paper)..... 175–200
Medicago sativa, silage compared with that from *Melilotus* spp..... 113–132
Mediterranean fruit fly. See *Ceratitis capitata*.
Melampsora—
lini, multinucleate fusion cells..... 646
pinitorqua, not cause of damping-off in conifers..... 550
Melanospora sp., fungus isolated from *Ipomoea batatas*..... 364
Melicope triphylla, susceptibility to *Pseudomonas citri*..... 663–665
Melilotus spp., silage compared with that from *Medicago sativa*..... 113–132
Melon fly. See *Bactrocera cucurbitae*.
Melon—
pomegranate. See *Cucumis melo* var. *dudaim*.
snake. See *Cucumis melo*.
Mercuric— Page
chlorid—
effect on—
Bacterium lachrymans..... 205
oxidase activity of seeds..... 168–169
used in determining nitrogen in spinach. 399–404
nitrate, effect on oxidase activity of seeds.. 169
Mercurous nitrate, effect on oxidase activity of seeds..... 169
Merrill, T. C., and Hartley, Carl (paper): Seedling Diseases of Conifers..... 521–558
Metabolism, nitrogen, in normal and blighted spinach..... 385–404
Meta-cresol, used to determine oxidase activity in normal and blighted spinach... 377–380
Methane fermentation in alfalfa hay and starch..... 274–275
Microcitrus—
australasica, susceptibility to *Pseudomonas citri*..... 662–665
australis, susceptibility to *Pseudomonas citri*..... 662–665
Microbial flora of alfalfa silage..... 574–592
Mimusops elengi, used in obtaining oviposition records for *Opius humilis*..... 443
Miris—
abbreviatus. See *Miris dolabratus*.
belangeri. See *Miris dolabratus*.
dolabratus—
control measures..... 197–198
description..... 191–194
dimorphism..... 183–184
distribution..... 175–181
economic importance..... 181–182
food plants..... 182–183
life history..... 184–194
natural enemies..... 194–197
synonymy..... 183
lateralis. See *Miris dolabratus*.
picticeps. See *Miris dolabratus*.
Moisture—
effect on—
after-ripening of seeds..... 138
soil acidity..... 321–329
toxity of alkali..... 306–309
in healthy and blighted spinach..... 387–390
in sweet-clover and alfalfa silage..... 114
Molasses, effect on alfalfa silage..... 574–590
Momordica—
balsamina, free from attacks of *Bacterium lachrymans*..... 206
charantia, free from attacks of *Bacterium lachrymans*..... 206
elaterium, free from attacks of *Bacterium lachrymans*..... 206
Monilochaetes infuscans, causal organism of scurf in *Ipomoea batatas*..... 352
Monoamino acid nitrogen in healthy and blighted spinach..... 399–404
Monosaccharids in hardened and nonhardened cabbage..... 95–96
Moss roses, resistant to brown canker..... 593–600
Moth—
brown-tail. See *Euproctis chrysorrhoea*.
gypsy. See *Porthetria dispar*.
rusty vapor. See *Notolophus antiquus*.

Mucor— Page
racemosus, fungus isolated from *Ipomoea batatas* ... 354-355
sp., not cause of damping-off in conifers ... 549
Multiple-Pipette Holder for the Distribution of Serum for the Complement-Fixation Test, A (paper) ... 615-618
Murraya exotica. Syn. *Chalcas exotica*.
Muskmelon. See *Cucumis melo*.
Nectria ipomoeae, fungus isolated from *Ipomoea batatas* ... 363-364
Neocosmospora vasinfectum, fungus isolated from *Ipomoea batatas* ... 364
Neofabraea malicorticis, fungus used in inoculation experiments with *Ipomoea batatas* ... 362
Net Energy Values of Alfalfa Hay and of Starch (paper) ... 269-286
Newsom, L. E., and Glover, George H. (paper): Further Studies on Brisket Disease ... 409-414
Nitrate—
calcium, in solution used in experiments with greensand ... 486-492
mercuric, effect on oxidase activity of seeds. 169
mercurous, effect on oxidase activity of seeds ... 169
potassium—
method, used in determining soil acidity ... 322-329
used in experiments with greensand ... 488-492
silver, used in determining nitrogen in spinach ... 399-404
sodium, effect on plowsole ... 507
Nitric acid. See Acid, nitric.
Nitrifying organisms in soil, effect of carbon disulphid and toluol ... 601-614
Nitrogen—
acid amid, in healthy and blighted spinach ... 398-404
amino, in alfalfa silage ... 581-590
ammonia, in healthy and blighted spinach ... 398-404
basic, in healthy and blighted spinach ... 399-404
digestibility in alfalfa hay and starch ... 271, 282
humin, in healthy and blighted spinach. 399-404
in dry matter of feeding stuffs ... 270
in hardened and nonhardened cabbage. 97, 105-108
in silage extract ... 125-132
metabolism in normal and blighted spinach ... 385-404
monoamino acid, in healthy and blighted spinach ... 399-404
peptid, in normal and blighted spinach. 400-404
residual soluble, in normal and blighted spinach ... 400-404
Nitrogen-fixing organisms in soil, effect of carbon disulphid and toluol ... 601-614
Nitrogen-free extract, digestibility in alfalfa hay and starch ... 270, 271, 282
Nitrohydrochloric acid. See Acid, nitrohydrochloric.
Nonprotein content of dry matter of feeding stuffs ... 270
Nonprotein nitrogen in normal and blighted spinach ... 393-397
Nonproteins in normal and blighted spinach 385-404
Norway spruce. See *Picea excelsa*.

Page
Notolophus antiquus, used in experiments with *Limnerium validum* ... 458
Nuts, kamani. See *Terminalia catappa*.
Oats. See *Avena sativa*.
Observations and Experiments on Intestinal Trichinæ (paper) ... 467-482
Observations on an Outbreak of Favus (paper) ... 415-418
Olive earth, overlying greensand ... 484
Olive, Russian wild. See *Elaeagnus* sp.
Oncognathus binotatus, infesting timothy .. 178, 181
Oomycetes causing damping-off of conifers ... 530
Opiinae, three species natural enemies of Mediterranean fruit fly ... 419-466
Opius—
fletcheri—
enemy of Mediterranean fruit fly ... 419-466
melon-fly parasite ... 456
humilis—
life history ... 440-445
natural enemy of Mediterranean fruit fly ... 419-466
Orchard grass. See *Dactylis glomerata*.
Orchids, cattleya, fumigation with hydrocyanic-acid gas ... 263-268
Organic matter—
digestibility in alfalfa hay and starch ... 271, 282
effect on toxity of alkali ... 303-305
Orthoptera, found on imported orchids ... 263
Osborn, Herbert (paper): The Meadow Plant Bug, Miris dolabratus ... 175-200
Oxid—
aluminium, in—
mulch, plowsole, and subsoil ... 508-516
normal and blighted spinach ... 371-375
soil, effect of drying ... 325-328
calcium, in—
greensand ... 485
mulch, plowsole, and subsoil ... 515-516
normal and blighted spinach ... 371-375
soil, effect of drying ... 325-328
cuprous, in intumescences on frozen leaves. 88
ferric, in—
mulch, plowsole, and subsoil ... 508-516
soil, effect of drying ... 325-328
magnesium, in—
normal and blighted spinach ... 371-375
soil, effect of drying ... 325-328
manganese, in soil, effect of drying ... 325-328
manganous, in mulch, plowsole, and subsoil ... 515-516
potassium, in—
greensand ... 485
normal and blighted spinach ... 371-375
red manganese, in normal and blighted spinach ... 371-375
silicic, in soil, effect of drying ... 325-328
sodium, in normal and blighted spinach. 371-375
Oxidase, in—
frozen cabbage leaves ... 88
nonliving structures of seeds ... 167-168
normal and blighted spinach ... 377-380
plant tissues, relation to manganese content 373
seeds ... 137-174
Oxygen—
effect on after-ripening of seeds ... 138
liberated from seeds by catalase activity. 142-161

Page

Pachycrepoideus dubius—
enemy of Mediterranean fruit fly..... 419–466
puparium parasite of Mediterranean fruit fly..... 461–463
Panicularia pauciflora, source of *Puccinia graminis avenae* transferred to uncongenial hosts..... 243
Panicum capillare, food plant of *Miris dolabratus*..... 182
Paracresol, used to determine—
catalase activity of seeds..... 162, 166
oxidase activity in normal and blighted spinach..... 377–380
Paramignya longipedunculata, susceptibility to *Pseudomonas citri*..... 662–665
Parasites of Mediterranean fruit fly in Hawaii..... 419–466
Parasitism, Morphology, and Cytology of Cronartium ribicola (paper)..... 619–660
Peach. See *Amygdalus persica*.
Pemberton, C. E., and Willard, H. F. (paper): Contribution to the Biology of Fruit-Fly Parasites in Hawaii..... 419–466
Penicillium—
sp., fungus isolated from *Ipomoea batatas*..... 355–356, 364
sp., not cause of damping-off in conifers..... 549
Pentoxid, phosphorus, in—
greensand..... 485
mulch, plowsole, and subsoil..... 508–516
normal and blighted spinach..... 371–375
Peonia officinalis, host of *Cronartium flaccidum*..... 638–639
Peptid nitrogen in normal and blighted spinach..... 400–404
Peridermium—
cerebrum, parasite of *Pinus virginiana*..... 625
pini—
parasite on *Pinus sylvestris*..... 625
acicolum, nuclear division of æciospores.. 639
Peroxid of hydrogen, used in determining catalase content of seeds..... 140–144
Peroxidase in frozen cabbage leaves..... 88–89
Pestalozzia—
funerea, isolated from damped-off conifers 547–558
hartigii, isolated from damped-off conifers. 547–558
sp., isolated from *Ipomoea batatas*..... 364
Pheidole megacephala, natural enemy of Mediterranean fruit fly..... 419–466
Phenolphthalein, used in—
determining catalase content of seeds..... 139
titration of silage extract..... 116–117
Phleum pratense—
food plant of *Miris dolabratus*..... 176, 182–183
source of *Puccinia graminis phleipratensis* transferred to semicongenial hosts..... 244
uncongenial host of *Puccinia graminis avenae*..... 243
Phloroglucin, used to determine oxidase reaction in normal and blighted spinach.. 377–380
Phoma—
batatae, reported causal organism of dryrot of *Ipomoea batatas*..... 349
betae, not cause of damping-off in conifers. 540–558
Phorantha occidentis, parasite of *Miris dolabratus*..... 176

Phosphate— Page
potassium, in solution used in experiments with greensand..... 486–492
sodium, effect on plowsole..... 507
Phosphates in—
greensand..... 483–492
olive earth..... 484
Phosphomolybdic acid. See Acid, phosphomolybdic.
Phosphorus—
in hardened and nonhardened cabbage..... 97, 105–108
pentoxid in—
greensand..... 485
mulch, plowsole, and subsoil..... 508–516
normal and blighted spinach..... 371–375
Phosphotungstate, barium, used in determining nitrogen in spinach..... 399–404
Phosphotungstic acid. See Acid, phosphotungstic.
Physiological Studies of Normal and Blighted Spinach (paper)..... 369–408
Phytophthora sp., causal organism of damping-off in conifers..... 530
Picea—
engelmanni, host of damping-off parasites. 523–558
excelsa, host of damping-off parasites..... 531–558
Piemeisel, F. J., et al. (paper): Plasticity of Biologic Forms of Puccinia graminis..... 221–249
Pig, guinea, weight of heart..... 411
Pine—
Austrian. *Pinus austriaca*.
Corsican. See *Pinus nigra poiretiana*.
jack. See *Pinus banksiana*.
loblolly. See *Pinus taeda*.
pitch. See *Pinus rigida*.
red. See *Pinus resinosa*.
Scotch. See *Pinus sylvestris*.
scrub. See *Pinus virginiana*.
shortleaf. See *Pinus echinata*.
white. See *Pinus strobus*.
yellow. See *Pinus ponderosa*.
Pinus—
austriaca, host of damping-off parasites.. 531–558
banksiana, host of damping-off parasites. 525–558
echinata, germination in mineral soil..... 526
flexilis, host of *Cronartium ribicola*..... 620
montana, attacked by *Melampsora pinitorqua*..... 551
nigra poiretiana, host of damping-off fungi..... 524–558
parviflora—
host of *Cronartium ribicola*..... 620
nature of infection by *Cronartium ribicola*..... 649, 651
ponderosa—
host of damping-off parasites..... 523–558
nature of injury from *Cronartium comptoniae*..... 649
resinosa, host of damping-off parasites.... 531–558
rigida, germination in mineral soil..... 526
scopulorum. Syn. *Pinus ponderosa*.
strobus—
host of *Cronartium ribicola*..... 619–660
host of damping-off parasites..... 530–558
sylvestris, host of—
damping-off parasites..... 531–558
Peridermium pini..... 625

Pinus—Continued. Page
taeda, germination in mineral soil..... 526
virginiana—
germination in mineral soil..... 526
host of *Peridermium cerebrum*..... 625
Pipette holder for distribution of serum... 615-618
Pitch pine. See *Pinus rigida*.
Pittman, D. W., and Harris, F. S. (paper): Soil Factors Affecting the Toxity of Alkali..... 287-319
Plant bug—
four-lined. See *Poecilocapsus lineatus*.
meadow. See *Miris dolabratus*.
tarnished. See *Lygus pratensis*.
Plants, hardening against frost..... 83-111
Plasmolysis of cells—
caused by freezing..... 90-92
effect on hydrogen-ion concentration of sap. 99
Plasticity of Biologic Forms of Puccinia graminis (paper)..... 221-249
Plenodomus destruens, causal organism of foot-rot of *Ipomoea batatas*..... 350
Plot tests, field technic..... 251-262
Plowsole, relation of inorganic soil colloids to..... 505-519
Poa pratensis—
catalase and oxidase content of seeds.... 137-174
food plant of *Miris dolabratus*..... 182
Poecilocapsus lineatus, pest of currants..... 181
Polypeptids in normal and blighted spinach..... 385-404
Polyphagus euglenae, similarity to *Cronartium ribicola*..... 648
Polysaccharids in hardened and nonhardened cabbage..... 95-96
Pomegranate melon. See *Cucumis melo* var. *dudaim*.
Popillia japonica, larvæ, effect of hydrocyanic-acid gas on..... 134-136
Populus—
alba, attacked by *Melampsora pinitorqua*.. 551
balsamifera candicans, attached by *Melampsora pinitorqua*..... 551
tremula, attached by *Melampsora pinitorqua*..... 551
Porthetria dispar, used in experiments with *Limnerium validum*..... 458
Potash in—
greensand..... 483-492
Hagerstown silty loam, soluble in—
ammonium-chlorid solution..... 67-70
carbonated water..... 67
distilled water..... 64-66
hydrochloric acid..... 63-64
Potassium—
chlorid, used in experiments with greensand..... 488
cyanid, tolerance of larvæ of *Trichinella spiralis* to..... 477-479
in greensand..... 483-492
nitrate, used in experiments with greensand..... 488-492
oxid in—
normal and blighted spinach..... 371-375
greensand..... 485
phosphate in solution used in experiments with greensand..... 486-492
Potassium-alum solution, used in determining protein nitrogen in spinach..... 391-393
Potassium-nitrate method, used in determining soil acidity..... 322-329
Potato—
sweet. See *Ipomoea batatas*.
See *Solanum tuberosum*.
Potentiometric determination of changes in hydrogen-ion concentration of plant sap during freezing..... 99-101
Protein—
digestibility in alfalfa hay and starch.... 271, 287
in dry matter of feeding stuffs..... 270
in hardened and nonhardened cabbage. 97, 103-104
in normal and blighted spinach..... 385-404
Pseudomonas—
citri—
susceptibility of rutaceous plants..... 661-666
virulence to Rutaceae..... 661-666
lachrymans. Syn. *Bacterium lachrymans*.
Puccinia—
dispersa, plasticity of biologic forms..... 221
falcariae, æciospore chains..... 647
graminis—
plasticity of biologic forms..... 221-249
agrostis, plasticity of biologic forms.... 241-247
avenae, used in bridging-host experiments..... 227-249
phleipratensis, plasticity of biologic forms..... 241-247
secalis, used in bridging-host experiments..... 225
tritici, used in bridging-host experiments..... 225-249
tritici-compacti, used in bridging-host experiments..... 227-249
Pucciniastrum agrimoniae, structure of uredinium..... 652
Pulverization of seeds, effect on catalase activity..... 143-145
Pumpkin. See *Cucurbita pepo*.
Pyrocatechin, used as test for oxidase in frozen cabbage leaves..... 88
Pyrocatechol, used to determine oxidase activity of—
normal and blighted spinach..... 377-380
seeds..... 162, 166
Pyrogallol, used to determine oxidase in—
frozen cabbage leaves..... 88
normal and blighted spinach..... 377-380
seeds..... 162, 166
Pseudotsuga taxifolia, host of damping-off parasites..... 531-558
Pythium debaryanum, cause of damping-off in conifers..... 522-558
Rabbit, weight of heart..... 411
"Rain-barley," relation to after-ripening..... 153
Rambler roses, resistant to brown canker.. 593-600
Reagents used in determining oxidase activity of seeds..... 162, 166
Red clover. See *Trifolium pratense*.
Red manganese oxid in normal and blighted spinach..... 371-375
Red pine. See *Pinus resinosa*.
Reducing sugars in—
hardened and nonhardened cabbage..... 95-96
healthy and blighted spinach..... 381-384
Reduviolus ferus, natural enemy of *Miris dolabratus*..... 194, 196

Page

Relation of Inorganic Soil Colloids to Plowsole in Citrus Groves in Southern California 505-519
Residual soluble nitrogen in normal and blighted spinach 400-404
Respiration of seeds, relation of catalase and oxidase content to 137-174
Reynolds, Francois H. (paper): A Multiple-Pipette Holder for the Distribution of Serum for the Complement-Fixation Test 615-618
Rheosporangium aphanidermatus, causal organism of damping-off in conifers 530, 540
Rhizoctonia, causal organisms of damping-off in conifers 522-558
Rhizoctonia—
crocorum, not grown on artificial media ... 530
medicaginis. Syn., *Rhizoctonia crocorum*.
Rhizopus nigricans, causal organism of sweet-potato storage-rots 338-368
Ribes spp., hosts of *Cronartium ribicola* 619-660
Ringrot of *Ipomoea batatas*—
dissemination 342
inoculation experiments 342-344
symptoms 340-342
Rosa spp., resistant to brown canker 593-600
Coniothyrium fuckelii, difference from *Diaporthe umbrina* 593-594
Rosellinia sp., not cause of damping-off in conifers 556
Roses. See *Rosa* spp.
Rugosas, resistant to brown canker 593-600
Russian wild olive. See *Elaeagnus* sp.
Rust, timothy, transferred by bridging hosts. 222
Rusty vaporer moth. See *Nololophus antiquus*.
Rutaceous plants, susceptibility to citrus canker 661-666
Rye—
effect on alfalfa silage 574-590
See *Secale cereale*.
Salt content of soils, determination by freezing-point method 331-336
Salvia, production of injected areas due to freezing 86
Sanford, H. L., and Sasscer, E. R. (paper): Effect of Hydrocyanic-Acid Gas under Vacuum Conditions on Subterranean Larvæ 133-136
Sasscer, E. R., and Dietz, H. F. (paper): Fumigation of Cattleya Orchids with Hydrocyanic-Acid Gas 263-268
Sasscer, E. R., and Sanford, H. L. (paper): Effect of Hydrocyanic-Acid Gas under Vacuum Conditions on Subterranean Larvæ 133-136
Schwartz, Benjamin (paper): Observations and Experiments on Intestinal Trichinæ. 467-482
Sclerotium—
bataticola, causal organism of charcoal-rot in *Ipomoea batatas* 351
rolfsii, fungus isolated from *Ipomoea batatas*. 364
Scofield, C. S. (paper): Effect of Farm Manure in Stimulating the Yields of Irrigated Farm Crops 493-503
Scotch pine. See *Pinus sylvestris*.
Scrub pine. See *Pinus virginiana*.
Scurf, fungous disease of *Ipomoea batatas* ... 352-353

Page

Secale cereale, used in bridging-host experiments 226-249
Seed, dormant, decay due to damping-off 525
Seedling Diseases of Conifers (paper) 521-558
Seedlings, conifer, injury from—
damping-off 521-558
drouth 554-555
mechanical means 551-554
whitespot 551
Seeds—
catalase and oxidase content of 137-174
effect of—
after-ripening on catalase activity 152-156
after-ripening on oxidase activity 165-166
age on catalase activity 150-154
age on oxidase activity 164-165
bromelin on oxidase activity 169
drying on catalase activity 149-150
germination on oxidase activity 166-167
heat on catalase activity 154-156
maturity on catalase activity 148-149
maturity on oxidase activity 163-164
mercuric chlorid on oxidase activity ... 168-169
mercuric nitrate on oxidase activity 169
mercurous nitrate on oxidase activity 169
retention in germinator on catalase activity 156-159
vitality on catalase activity 150-154
oxidase activity of nonliving structures .. 167-168
Serum, distributed by multiple-pipette holder 615-618
Severinia buxifolia, immunity to *Pseudomonas citri* 662-665
Shortleaf pine. See *Pinus echinata*.
Silage—
alfalfa—
acidity 374, 581
bacteria in 571-592
bacteriological analysis 586-587
chemical analysis 586-587
effect of carbohydrates 584-590
loss in weight 114
moisture content 114
sweet-clover compared with alfalfa 113-132
Silica in—
greensand 483-492
mulch, plowsole, and subsoil 508-516
Silicic-oxid content of soil, effect of drying. 325-328
Silicon dioxid in—
mulch, plowsole, and subsoil 508-516
normal and blighted spinach 371-375
Silver—
chlorid, used in determining nitrogen in spinach 400-404
nitrate, used in determining nitrogen in spinach 399-404
Snake melon. See *Cucumis melo*.
Sodium—
carbonate—
effect on plowsole 507
used in alkali experiment 289-319
chlorid—
effect on plowsole 507
used in alkali experiment 289-319
used in experiments with intestinal trichinæ 468-482

Sodium—Continued. Page
cyanid—
effect on cattleya orchids ... 267–268
used to generate hydrocyanic-acid gas ... 134
hydroxid—
in alfalfa silage ... 586–590
used to determine—
catalase content of seeds ... 139
water-soluble nitrogen in spinach ... 398–404
nitrate, effect on plowsole ... 507
oxid in normal and blighted spinach ... 371–375
phosphate, effect on plowsole ... 507
sulphate, used in alkali experiment ... 289–319
Softrot in *Ipomoea batatas*—
dissemination ... 342
inoculation experiments ... 342–344
symptoms ... 340–342
Soil Acidity as Affected by Moisture Conditions of the Soil (paper) ... 321–329
Soil—
adobe, inorganic colloids ... 516
colloids, relation to plowsole ... 505–519
effect of carbon disulphid and toluol on nitrogen-fixing organisms ... 601–614
Hagerstown silty loam, fertilizer potash residues in ... 59–81
overwintering of *Bacterium lachrymans* in ... 213
particles, effect of size on toxity of alkali ... 290–301
salt content of, determination by freezing-point method ... 331–336
water content of, effect on fumigation with hydrocyanic-acid gas ... 133–136
Soil Factors Affecting the Toxicity of Alkali (paper) ... 287–319
Solanum tuberosum—
effect of farm manure on ... 494–503
leafroll ... 559–570
Sorghum halepensis—
diameter of cells in seeds ... 145
catalase and oxidase content of seeds ... 137–174
hybridized with Tunis grass, catalase activity of seeds ... 146–147
Species, new ... 593–600
Sphaeronema fimbriatum, causal organism of blackrot of *Ipomoea batatas* ... 344
Spinacia oleracea—
denitrification due to blight ... 401–404
effect of blight on—
ash content ... 371–375
carbohydrate production ... 381–384
nitrogen metabolism ... 385–404
oxidase reaction ... 377–380
Spinach-blight, physiological studies ... 369–408
Sporobolus cryptandrus, source of *Puccinia graminis* in bridging-host experiment ... 228
Sporotrichium sp., fungus isolated from *Ipomoea batatas* ... 364
Spruce—
Engelmann. See *Picea engelmanni*.
Norway. See *Picea excelsa*.
Squash—
Hubbard. See *Cucurbita maxima*.
Japanese crookneck. See *Cucurbita moschata*.
summer. See *Cucurbita pepo* var. *condensa*.
Squirting cucumber. See *Momordica elaterium*.

Page
Stakman, E. C., Piemeisel, F. J., and Levine, M. N. (paper): Plasticity of Biologic Forms of Puccinia graminis ... 221–249
Starch—
digestibility ... 271, 282
influence on digestibility of hay ... 272
in hardened and nonhardened cabbage ... 95–96, 105–108
in healthy and blighted spinach ... 381–384
methane fermentation ... 274–275
net energy value ... 269–286
Storage, effect on oxidase activity of seeds ... 165–166
Storage-rots of *Ipomoea batatas* ... 337–368
Straw, effect on alfalfa silage ... 574–590
Studies in Inheritance of Certain Characters of Crosses between Dairy and Beef Breeds of Cattle (paper) ... 1–57
Stutzer's nitrogen in silage ... 129–132
Sucrose in—
hardened and nonhardened cabbage ... 95–96
healthy and blighted spinach ... 381–384
Sudan grass. See *Andropogon halepensis sudanensis*.
Sugar beets. See *Beta vulgaris*.
Sugar, cane, effect on alfalfa silage ... 584–590
Sugars in healthy and blighted spinach ... 381–384
Sulphate—
calcium, effect on plowsole ... 507
copper, effect on *Bacterium lachrymans* ... 205
magnesium—
effect on plowsole ... 507
in solution used in experiments with greensand ... 486–492
sodium, used in alkali experiment ... 289–319
Sulphid, copper, used in determining nitrogen in spinach ... 400–404
Sulphur trioxid in—
greensand ... 485
normal and blighted spinach ... 371–375
Sulphuric acid. See Acid, sulphuric.
Sunflower. See *Helianthus annuus*.
Swanson, C. O., and Tague, E. L. (paper): Chemistry of Sweet-Clover Silage in Comparison with Alfalfa Silage ... 113–132
Sweet clover. See *Melilotus* spp.
Sweet-clover silage compared with alfalfa silage ... 113–132
Sweet-Potato Storage-Rots (paper) ... 337–368
Tague, E. L., and Swanson, C. O. (paper): Chemistry of Sweet-Clover Silage in Comparison with Alfalfa Silage ... 113–132
Tarnished plant bug. See *Lygus pratensis*.
Taxus—
canadensis, host of damping-off parasites ... 531–558
cuspidata, host of damping-off parasites ... 531–558
Temperature—
effect on—
Bacterium lachrymans ... 204–205
catalase activity of seeds ... 154–156
relation to hardening of plants against frost injury ... 93–94
Terminalia catappa, host of puparia parasitized by *Diachasma tryoni* ... 429
Tetramethylparaphenylenediamin, used as a test for oxidase in frozen cabbage leaves ... 88

Tetrastichus giffardianus— Page
life history ... 448-454
natural enemy of Mediterranean fruit fly. 419-466
Thevetia neriifolia—
host of puparia parasitized by *Diachasma fullawayi* ... 448
host of larvæ parasitized by *Tetrastichus giffardianus* ... 452
Thymolphthalein, used in titration of silage extract ... 119-127
Tilia americana, catalase and oxidase content of seeds ... 138
Timothy—
rust, transferred by bridging hosts ... 222
See *Phleum pratense*.
Titration of silage extract ... 116-130
Toddalia asiatica, susceptibility to *Pseudomonas citri* ... 663-665
Toluol—
effect on nitrogen-fixing organisms in soil 601-614
used in determining carbohydrates in normal and blighted spinach ... 383
Tomato. See *Lycopersicon esculentum*.
Toxity of alkali, effect of soil factors ... 287-315
Trichimella spiralis, growth and molting ... 467-482
Trichoderma—
köningi, isolated from—
damped-off conifers ... 546-558
Ipomoea batatas ... 360-361.
lignorum—
isolated from damped-off conifers ... 539-558
used in inoculation experiments with *Ipomoea batatas* ... 361
Trichosanthes colubrina, free from attacks of *Bacterium lachrymans* ... 206
Trichosporium sp., fungus isolated from *Ipomoea batatas* ... 364
Trichothecium roseum, not cause of damping-off in conifers ... 540-558
Trifolium—
incarnatum—
catalase content of seeds ... 144
diameter of cells in seeds ... 145
pratense, used in experiments with greensand ... 485-492
Trioxid, sulphur, in—
greensand ... 485
normal and blighted spinach ... 371-375
Triphasia trifolia, immunity to *Pseudomonas citri* ... 662-665
Triticum—
aestivum—
catalase activity of parts of seed ... 146
used in bridging-host experiments ... 226-249
used in experiments with greensand ... 485-492
yield on alkali land ... 302
vulgare, reported infected with timothy rust through bridging host ... 222

Page
True, Rodney H., Black, Otis F., Kelly, James W., Bunzell, H. H., Hawkins, Lon A., Jodidi, Samuel L., and Kellogg, Edward H. (paper): Physiological Studies of Normal and Blighted Spinach ... 369-408
True, Rodney H., and Geise, Fred W. (paper): Experiments on the Value of Greensand as a Source of Potassium for Plant Culture ... 483-492
Tunis grass and sorghum hybrid, catalase activity of seeds ... 146-147
Turban squash. *Cucurbita pepo* var. *condensa*.
Tyrosin—
used to determine oxidase activity in normal and blighted spinach ... 377-380
test for oxidase in frozen cabbage leaves ... 83
Tyrosine, oxidase of Johnson grass not active toward ... 162
Urinary excretion, analysis in alfalfa hay and starch-feeding experiment ... 273, 283
Vacuum process of fumigation with hydrocyanic-acid gas ... 133-136
Vaseline, used in treatment of favus ... 417-418
Verticillium cinnabrinus, fungus isolated from *Ipomoea batatas* ... 364
Vitality of seeds, relation of catalase and oxidase content to ... 137-174
Water in soil, effect on fumigation with hydrocyanic-acid gas ... 133-136
Water extract of silage—
acidity ... 116-125
method of making ... 114-115
nitrogen in ... 125-132
Watermelon. See *Citrullus vulgaris*.
Wax on plant leaves, effect upon freezing point of plant ... 86
Webworm, fall. See *Hyphantria cunea*.
Weimer, J. L., et al. (paper): Sweet-Potato Storage-Rots ... 337-368
West Indian gherkin. See *Cucumis anguria*.
Wheat. See *Triticum aestivum*.
Wheat wireworm. See *Agriotes mancus*.
White grub. See *Lachnosterna* sp.
White pine. See *Pinus strobus*.
Whitespot, resemblance to damping-off in conifer seedlings ... 551
Wild cucumber. See *Echinocystis lobata*.
Willard, H. F., and Pemberton, C. E. (paper): Contribution to the Biology of Fruit-Fly Parasites in Hawaii ... 419-466
Wireworm, wheat. See *Agriotes mancus*.
Witch grass. See *Panicum capillare*.
Xanthoxylum rhetsa, immunity to *Pseudomonas citri* ... 663-665
Yeasts in alfalfa silage ... 574-592
Yellow pine. See *Pinus ponderosa*.
Yew. See *Taxus cuspidata*.
Zygorhynchus sp., fungus isolated from *Ipomoea batatas* ... 364
Zymase, cause of alcohol in silage ... 590